

# Laterodorsal tegmental nucleus

From Wikipedia, the free encyclopedia

The **laterodorsal tegmental nucleus** (or

	<b>Brain: Laterodorsal tegmental nucleus</b>
<b>Latin</b>	<i>nucleus tegmentalis posterolateralis</i>
<b>NeuroNames</b>	<i>ancil-282</i> ( <i>http://braininfo.rprc.washington.edu/Scripts/ancilcentraldirectory.aspx?ID=282</i> )

**lateroposterior tegmental nucleus**) is a nucleus situated in the brainstem, spanning the midbrain tegmentum and the pontine tegmentum. Its location is one-third of the way from the pedunculopontine nucleus to the thalamus, inferior to the pineal gland.

## Function

The laterodorsal tegmental nucleus (LDT) sends cholinergic (acetylcholine) projections to many subcortical and cortical structures, including the thalamus, hypothalamus, substantia nigra (dopamine neurons), ventral tegmental area (dopamine neurons), cortex (with bidirectional connections with the prefrontal cortex).

The laterodorsal tegmental nucleus may be involved in modulating sustained attention or in mediating alerting responses, and also in the generation of REM sleep (along with the pedunculopontine nucleus).

Retrieved from "http://en.wikipedia.org/w/index.php?title=Laterodorsal\_tegmental\_nucleus&oldid=467041708"

Categories: Neuroanatomy | Neuroanatomy stubs

- 
- This page was last modified on 21 December 2011 at 16:13.
  - Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.